

## Correspondence

### First survey of the pathogenic fungus *Batrachochytrium salamandrivorans* in wild and captive amphibians in the Czech Republic

VOJTECH BALÁŽ<sup>1,2</sup>, MILIČ SOLSKÝ<sup>3</sup>, DAVID LASTRA GONZÁLEZ<sup>3</sup>, BARBORA HAVLÍKOVÁ<sup>3</sup>, JUAN GALLEGU ZAMORANO<sup>3</sup>, CRISTINA GONZÁLEZ SEVILLEJA<sup>3</sup>, LAURA TORRENT<sup>3</sup> & JIŘÍ VOJAR<sup>3</sup>

<sup>1</sup>) Department of Ecology and Diseases of Game, Fish and Bees, Faculty of Veterinary Hygiene and Ecology, University of Veterinary and Pharmaceutical Sciences Brno, Palackého tř. 1/3, 612 42 Brno, Czech Republic

<sup>2</sup>) Department of Biology and Wildlife Diseases, Faculty of Veterinary Hygiene and Ecology, University of Veterinary and Pharmaceutical Sciences Brno, Palackého tř. 1/3, 612 42 Brno, Czech Republic

<sup>3</sup>) Department of Ecology, Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 165 21 Prague 6, Czech Republic

Corresponding author: Jiří VOJAR, e-mail: vojar@fzp.czu.cz

Manuscript received: 8 June 2017

Accepted: 27 July 2017 by JÖRN KÖHLER

The recently discovered fungal pathogen *Batrachochytrium salamandrivorans* (MARTEL et al. 2013) (hereinafter referred to as *Bsal*) has already received significant scientific and public attention (e.g., MARTEL et al. 2014, VAN ROOIJ et al. 2015, YAP et al. 2015, STEGEN et al. 2017). The *Bsal* epidemic has so far been limited to European newts and salamanders found in the wild (Belgium, Germany and the Netherlands: MARTEL et al. 2013, SPITZEN-VAN DER SLUIJS et al. 2016) and in captive populations (Germany: SABINO-PINTO et al. 2015; United Kingdom: CUNNINGHAM et al. 2015). In the Netherlands, *Bsal* is responsible for the near extinction of wild fire salamander (*Salamandra salamandra*) populations (SPITZEN-VAN DER SLUIJS et al. 2013).

The Bern Convention Standing Committee has therefore announced Recommendation No. 176 on the prevention and control of the *Bsal* chytrid fungus. According to this recommendation, European countries should adopt measures that include establishment of monitoring programmes to control the possible further spread of the disease, especially in areas of high risk (e.g., areas near disease outbreaks), and develop emergency action plans that will allow prompt responses in case of *Bsal* occurrence (Council of Europe 2015).

The Czech Republic is a country with relatively high caudate species diversity (SILLERO et al. 2014) and shares a western border with Germany, a country with previously proven *Bsal* occurrence (SABINO-PINTO et al. 2015, SPITZEN-VAN DER SLUIJS et al. 2016). The country, and especially the capital city of Prague, has an active and sizeable community of exotic pet keepers and pet shops, and large

exotic pet fairs take place on a regular basis (HAVLÍKOVÁ et al. 2015). Furthermore, Prague and its surroundings are known to harbour wild populations of at least four native caudate species: fire salamander, smooth newt (*Lissotriton vulgaris*), alpine newt (*Ichthyosaura alpestris*) and great crested newt (*Triturus cristatus*) (ŠŤASTNÝ et al. 2015). All four are susceptible to *Bsal*-induced mortality (MARTEL et al. 2013, 2014, CUNNINGHAM et al. 2015, SPITZEN-VAN DER SLUIJS et al. 2016, STEGEN et al. 2017). The surrounding areas of such large cities are likely to constitute areas of high risk for wild populations of native amphibians. For this reason, we selected Prague and its surroundings as the first focal area for *Bsal* surveillance efforts in wild populations of Czech caudate amphibians. Considering that *Bsal* is spread through the pet trade in caudates originating in Asia (MARTEL et al. 2014), we also focused on captive collections of caudate amphibians.

In total, 324 swab samples were tested for both *Batrachochytrium dendrobatidis* (*Bd*) and *Bsal* presence and prevalence. In wild populations, 126 samples of three caudate species (fire salamanders, smooth newts and alpine newts) were collected at nine sites within Prague's urban area and its surroundings during autumn 2015 and spring 2016 (Table 1). Furthermore, we analysed samples from five captive collections, including those of four private breeders and Prague's zoological garden during the period 2015–2016. Within each collection, only subset of about two to four individuals were sampled from an aquarium. This *Bsal*-targeted sampling in captivity was extended by re-analysing samples of caudates from previous surveillance projects

Table 1. Numbers (No.) of individuals sampled within nine wild caudate populations in Prague and its surroundings. Date = date of sampling.

Locality name	Coordinates		Species	No.	Date
Podhořský potok, small stream, tributary of Vltava River	50.129947°N	14.404111°E	<i>Salamandra salamandra</i>	31	07.10.2015
Únětice, unnamed tributary of Únětický potok	50.144853°N	14.384502°E	<i>Salamandra salamandra</i>	25	15.10.2015
Levý Hradec, unnamed tributary of Vltava River	50.169883°N	14.377429°E	<i>Salamandra salamandra</i>	12	20.10.2015
Úholičky, unnamed tributary of Podmoráňský potok	50.170698°N	14.344784°E	<i>Salamandra salamandra</i>	8	09.11.2015
Lhotecký potok, tributary of Vltava River	49.956059°N	14.411423°E	<i>Salamandra salamandra</i>	7	15.10.2015
Chalupecká strouha, near confluence with Zvolský potok	49.930541°N	14.390361°E	<i>Salamandra salamandra</i>	1	17.11.2015
Baně, unnamed tributary of Vltava River	49.961229°N	14.392828°E	<i>Salamandra salamandra</i>	2	17.11.2015
Ohrobecké údolí, unnamed tributary of Vltava River	49.943775°N	14.413338°E	<i>Salamandra salamandra</i>	10	21.10.2015
Botanická zahrada, artificial pond in botanic garden	50.070429°N	14.421077°E	<i>Lissotriton vulgaris</i>	28	01.07.2016
Botanická zahrada, artificial pond in botanic garden			<i>Ichthyosaura alpestris</i>	2	01.07.2016

searching for *Bd* presence in captive amphibians (HAVLÍKOVÁ et al. 2015), including 18 individuals of the largest amphibian species, the Chinese giant salamander (*Andrias davidianus*), reared in Prague's zoological garden. In total, 198 samples of 60 caudate (sub)species were analysed in captive collections (Table 2).

Sampling and DNA extraction were performed according to procedures used in amphibian chytridiomycosis research (BOYLE et al. 2004). The first sample subset, consisting of 98 wild and 56 captive samples, was checked for *Bsal* presence by SYBRGreen quantitative polymerase chain reaction (qPCR) following the method described in BLOOI et al. (2013) as one possible detection option. Bovine serum albumin (BSA) was added to reduce PCR inhibition (GARLAND et al. 2010). The identity of amplified DNA was checked by melt curve analysis and compared to results for genomic standards of *Bsal* provided by An Martel (Ghent University). We later adopted the duplex *Bd+Bsal* qPCR (BLOOI et al. 2013) and used it for additional samples. In this assay, we used genomic standards of *Bd* equivalent to 0.1, 1, 10 and 100 zoospores per 5 µl (strain IA042, Ibon Acherito, Pyrenees, 2004) obtained from the Institute of Zoology, Zoological Society of London. A single quantity sample of *Bsal* genomic DNA was used as a positive control. If any sample showed fluorescence growth in the wavelength of the *Bsal* probe, it would be re-analysed with the full set of *Bsal* standards. In this way, we slightly reduced the cost of analysis. In both detection assays, we used duplicates of all analysed samples, standards, as well as positive and negative controls.

All tested samples yielded negative results for the presence of *Bsal*. *Bd* was detected in three individuals of wild smooth newts and in one reared ribbed newt (*Pleurodeles waltl*) in a captive collection, albeit with no visible signs of the chytridiomycosis. Low *Bd* prevalence in caudates corresponds well with our previous findings in Czech captive collections (HAVLÍKOVÁ et al. 2015), and wild populations of caudates in Central and east Europe (BALÁŽ et al. 2014a,b, VOJAR et al. 2017).

The 0% *Bsal* prevalence in wild caudates has Sterne-Wald 99% confidence limits of 0.0–4.2%, while in the case of sam-

ples from captivity the 99% confidence limits are 0.0–2.6% (RÓZSA et al. 2000). This does not directly mean that *Bsal* is not present in the Czech Republic. Because the disease outbreaks can occur at very low host densities in wild populations (SCHMIDT et al. 2017), all host populations of susceptible European caudate species (MARTEL et al. 2014) are at risk from *Bsal* (SCHMIDT et al. 2017). In the case of asymptomatic Asian caudates in captive collections, infection may be present in such small prevalence (MARTEL et al. 2014, LAKING et al. 2017) that our sampling was not sufficient. On the other hand, because the intensive sampling of wild fire salamanders covered nearly all localities within Prague where the species presently is known to occur (ŠTASTNÝ et al. 2015) and no sampled individual exhibited visible disease symptoms, we conclude that *Bsal* probably has not invaded Prague's fire salamander population, at least for now. Similar results of pathogen absence have been found in studies focused on fire salamanders in Austria (GIMENO et al. 2015), eastern hellbenders (*Cryptobranchus alleganiensis*) in the U.S. (BALES et al. 2015), Japanese giant salamanders (*Andrias japonicus*) in Japan (BLETZ et al. 2017a), Chinese amphibians (ZHU et al. 2014), five species of newts and fire salamanders in most of tested localities in Belgium, Germany and the Netherlands (SPITZEN-VAN DER SLUIJS et al. 2016), alpine newts, smooth newts and great crested newts in Germany (BLETZ et al. 2017b), and in a study by PARROT et al. (2016) on 17 caudate species across three continents.

We used two available detection assays in our study, both based on DNA amplification with the same pair of *Bsal* primers (BLOOI et al. 2013) and differing only in the detection format of the amplicon. The SYBR Green qPCR assay often produced detectable fluorescence growth of nonspecific products, thus complicating interpretation of the results. In several cases, we ran standard PCR followed by gel electrophoresis with samples of equivocal results to confirm the identity of PCR products. Our results indicated a mean melting temperature ( $T_m$ ) for *Bsal* standards of 77.21°C (SD = 0.29), which differs slightly from the published value of 75.5°C (BLOOI et al. 2013). For monitoring *Bsal* presence in wild and captive amphibians, we later adopted and recommend the use of duplex *Bd+Bsal* qPCR,

Table 2. List of surveyed species and numbers (No.) of individuals sampled in captivity.

Species	No.	Species	No.
<i>Ambystoma mexicanum</i>	3	<i>Neurergus deryugina deryugina</i>	2
<i>Ambystoma tigrinum</i>	2	<i>Neurergus strauchii barani</i>	3
<i>Andrias davidianus</i>	18	<i>Neurergus strauchii strauchii</i>	3
<i>Calotriton asper</i>	3	<i>Ommatotriton ophryticus nesterovi</i>	3
<i>Cynops ensicauda ensicauda</i>	3	<i>Pachyhynobius shangchengensis</i>	1
<i>Cynops ensicauda popei</i>	4	<i>Pachytriton</i> sp.	2
<i>Cynops orientalis</i>	1	<i>Paramesotriton caudopunctatus</i>	7
<i>Cynops pyrrhogaster</i>	3	<i>Paramesotriton deloustali</i>	6
<i>Cynops pyrrhogaster</i> “Kanagawa”	6	<i>Paramesotriton guangxiensis</i>	4
<i>Cynops pyrrhogaster</i> “Yubana”	2	<i>Paramesotriton hongkongensis</i>	3
<i>Euproctus platycephalus</i>	2	<i>Paramesotriton chinensis</i>	12
<i>Hynobius dunni</i>	1	<i>Paramesotriton</i> sp. “helm”	1
<i>Hynobius leechii</i>	2	<i>Paramesotriton</i> sp. “red”	6
<i>Hynobius lichenatus</i>	1	<i>Paramesotriton yunwensis</i>	2
<i>Hynobius quelpartensis</i>	2	<i>Pleurodeles nebulosus</i>	2
<i>Hynobius retardatus</i>	2	<i>Pleurodeles walzl</i>	4
<i>Hypselotriton cyanurus</i>	2	<i>Salamandra algira tingitana</i>	2
<i>Hypselotriton cyanurus cyanurus</i>	2	<i>Siren intermedia</i>	1
<i>Hypselotriton chenggongensis</i>	3	<i>Triturus anatolicus</i>	2
<i>Hypselotriton orientalis</i>	1	<i>Triturus blasii</i>	3
<i>Ichthyosaura alpestris</i>	3	<i>Triturus carnifex</i>	9
<i>Laotriton laoensis</i>	3	<i>Triturus cristatus</i>	2
<i>Lissotriton boscai</i>	3	<i>Triturus dobrogicus dobrogicus</i>	2
<i>Lissotriton graecus</i>	3	<i>Triturus dobrogicus macrosoma</i>	3
<i>Lissotriton helveticus</i>	3	<i>Triturus ivanbureschi</i>	1
<i>Lissotriton italicus</i>	3	<i>Triturus karelinii</i>	2
<i>Lissotriton malcani</i>	3	<i>Triturus macedonicus</i>	10
<i>Lissotriton meridionalis</i>	3	<i>Triturus marmoratus</i>	2
<i>Lissotriton montandoni</i>	3	<i>Triturus pygmaeus</i>	3
<i>Neurergus crocatus complex</i>	3	<i>Tylotriton shanjing</i>	2

which is designed specifically to detect either of the pathogen species (BLOOI et al. 2013). The higher cost of analyses versus using fluorescent probes is counterbalanced by clearer and more specific results.

As not only newts, but also infected anurans and even waterfowl via scales on their feet, may promote fungal spread over large spatial distances (STEGEN et al. 2017), the spread of this emerging pathogen is difficult to predict, and we can expect the distribution of *Bsal* to change considerably over time. The risk that new points of entry for *Bsal* into Europe will occur via the pet trade is a constant threat that can be alleviated by collaboration among pet owners, the pet trade, veterinary authorities, and conservationists (SABINO-PINTO et al. 2015). It is essential to prevent this pathogen entering the wild amphibian populations (CUNNINGHAM et al. 2015), because there is no effective method to reduce the impact of chytridiomycosis in the field (GARNER et al. 2016). Therefore, our next planned steps in the Czech Republic include establishment and issue of biosecurity guidelines for owners of caudates, providing *Bsal*

detection in captive collections of amphibians, forming a network of continuously monitored localities in proximity to larger cities, and preparing an action plan in case of *Bsal* occurrence in collaboration with the Nature Conservation Agency of the Czech Republic, the State Veterinary Authority, and the Czech Ministry of Environment.

#### Acknowledgements

We thank A. MARTEL from Ghent University for providing the genomic standards for *Bsal*, T. GARNER from the Institute of Zoology, Zoological Society of London, for providing the genomic standards for *Bd*, and G. A. KIRKING for useful comments on the manuscript. We also thank all those zookeepers, institutions, private amphibian breeders, and pet shop sellers who cooperated voluntarily and provided amphibians for sampling. The study was supported by the Czech Ministry of Environment and by the University of Life Sciences Prague (grant no. 20164245). The field work was performed on the basis of permission awarded by the Nature Conservation Agency of the Czech Republic.

## References

- BALÁŽ, V., J. VOJAR, P. CIVIŠ, M. ŠANDERA & R. ROZÍNEK (2014a): Chytridiomycosis risk among Central European amphibians based on surveillance data. – *Diseases of Aquatic Organisms*, **112**: 1–8.
- BALÁŽ, V., J. VÖRÖS, P. CIVIŠ, J. VOJAR, A. HETTYEY, E. SÓS, R. DANKOVICS, R. JEHL, D. G. CHRISTIANSEN, F. CLARE, M. C. FISHER, T. J. W. GARNER & J. BIELBY (2014b): Assessing risk and guidance on monitoring of *Batrachochytrium dendrobatidis* in Europe through identification of taxonomic selectivity of infection. – *Conservation Biology*, **28**: 213–223.
- BALES, E. K., O. J. HYMAN, A. H. LOUDON, R. N. HARRIS, G. LIPPS, E. CHAPMAN, K. ROBLEE, J. D. KLEOPFER & K. A. TERRELL (2015): Pathogenic chytrid fungus *Batrachochytrium dendrobatidis*, but not *B. salamandrivorans*, detected on eastern hellbenders. – *PLoS ONE*, **10**: e0116405.
- BLETZ, M. C., M. VENCES, J. SABINO-PINTO, Y. TAGUCHI, N. SHIMIZU, K. NISHIKAWA & A. KURABAYASHI (2017a): Cutaneous microbiota of the Japanese giant salamander (*Andrias japonicus*), a representative of an ancient amphibian clade. – *Hydrobiologia*, **795**: 153–167.
- BLETZ, M. C., R. G. B. PERL, B. T. C. BOBOWSKI, L. M. JAPKE, C. C. TEBBE, A. B. DOHRMANN, S. BHUJU, R. GEFFERS, M. JAREK & M. VENCES (2017b): Amphibian skin microbiota exhibits temporal variation in community structure but stability of predicted *Bd*-inhibitory function. – *The ISME Journal*, **11**: 1521–1534.
- BLOOI, M., F. PASMANS, J. E. LONGCORE, A. SPITZEN-VAN DER SLUIJS, F. VERCAMMEN & A. MARTEL (2013): Duplex Real-Time PCR for rapid simultaneous detection of *Batrachochytrium dendrobatidis* and *Batrachochytrium salamandrivorans* in amphibian samples. – *Journal of Clinical Microbiology*, **51**: 4173–4177.
- BOYLE, D. G., D. B. BOYLE, V. OLSEN, J. A. MORGAN & A. D. HYATT (2004): Rapid quantitative detection of chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples using real-time Taqman PCR assay. – *Diseases of Aquatic Organisms*, **60**: 141–148.
- Council of Europe (2015): Convention on the Conservation of European Wildlife and Natural Habitats, 35th meeting, 1–4 Dec. 2015, Strasbourg, France. – Recommendation No. 176 (2015) on the prevention and control of the *Batrachochytrium salamandrivorans* chytrid fungus. [accessed 23 March 2017]
- CUNNINGHAM, A. A., K. BECKMANN, M. PERKINS, L. FITZPATRICK, R. CROMIE, J. REDBOND, M. F. O'BRIEN, P. GHOSH, J. SHELTON & M. C. FISHER (2015): Emerging disease in UK amphibians. – *Veterinary Record*, **176**: 468.
- GARLAND, S., A. BAKER, A. D. PHILLOTT & L. F. SKERRATT (2010): BSA reduces inhibition in a TaqMan® assay for the detection of *Batrachochytrium dendrobatidis*. – *Diseases of Aquatic Organisms*, **92**: 113–116.
- GARNER, T. W. J., B. R. SCHMIDT, A. MARTEL, F. PASMANS, E. MUTHS, A. A. CUNNINGHAM, C. WELDON, M. C. FISHER & J. BOSCH (2016): Mitigating amphibian chytridiomycoses in nature. – *Philosophical Transactions of the Royal Society B*, **371**: 20160207.
- GIMENO, A., M. MEIKL, A. PITT, M. WINKLER & U. G. BERNINGER (2015): Testing of Fire Salamanders around Salzburg for *Batrachochytrium salamandrivorans* within a school project. – *Journal on Protected Mountain Areas Research*, **7**: 72–76.
- HAVLÍKOVÁ, B., J. VOJAR & V. BALÁŽ (2015): First systematic monitoring of *Batrachochytrium dendrobatidis* in collections of captive amphibians in the Czech Republic. – *Amphibia-Reptilia*, **36**: 27–35.
- LAKING, A. E., H. N. NGO, F. PASMANS, A. MARTEL & T. T. NGUYEN (2017): *Batrachochytrium salamandrivorans* is the predominant chytrid fungus in Vietnamese salamanders. – *Scientific Reports*, **7**: 44443.
- MARTEL, A., A. SPITZEN-VAN DER SLUIJS, M. BLOOI, W. BERT, R. DUCATELLE, M. C. FISHER, A. WOELTJES, W. BOSMAN, K. CHIERS, F. BOSSUYT & F. PASMANS (2013): *Batrachochytrium salamandrivorans* sp. nov. causes lethal chytridiomycosis in amphibians. – *Proceedings of the National Academy of Sciences of the U.S.A.*, **110**: 15325–15329.
- MARTEL, A., M. BLOOI, C. ADRIAENSEN, P. VAN ROOIJ, W. BEUKEMA, M. C. FISHER, R. A. FARRER, B. R. SCHMIDT, U. TOBLER, K. GOKA, K. R. LIPS, C. MULETZ, K. R. ZAMUDIO, J. BOSCH, S. LÖTTERS, E. WOMBWELL, T. W. J. GARNER, A. A. CUNNINGHAM, A. SPITZEN-VAN DER SLUIJS, S. SALVIDIO, R. DUCATELLE, K. NISHIKAWA, T. T. NGUYEN, J. E. KOLBY, I. VAN BOXC-LAER, F. BOSSUYT & F. PASMANS (2014): Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. – *Science*, **346** (6209): 630–631.
- PARROTT, J. C., A. SHEPACK, D. BURKART, B. LABUMBARD, P. SCIME, E. BARUCH & A. CATENAZZI (2016): Survey of pathogenic chytrid fungi (*Batrachochytrium dendrobatidis* and *B. salamandrivorans*) in salamanders from three mountain ranges in Europe and the Americas. – *Ecohealth*, **14**: 296–302.
- RÓZSA, L., J. REICZIGEL & G. MAJOROS (2000): Quantifying parasites in samples of hosts. – *Journal of Parasitology*, **86**: 228–232.
- SABINO-PINTO, J., M. BLETZ, R. HENDRIX, R. G. B. PERL, A. MARTEL, F. PASMANS, S. LÖTTERS, F. MUTSCHMANN, B. R. SCHMIDT, D. S. SCHMELLER, M. VEITH, N. WAGNER, M. VENCES & S. STEINFARTZ (2015): First detection of the emerging fungal pathogen *Batrachochytrium salamandrivorans* in Germany. – *Amphibia-Reptilia*, **36**: 411–416.
- SCHMIDT, B. R., C. BOZZUTO, S. LÖTTERS & S. STEINFARTZ (2017): Dynamics of host populations affected by the emerging fungal pathogen *Batrachochytrium salamandrivorans*. – *Royal Society Open Science*, **4**: 160801.
- SILLERO, N., J. CAMPOS, A. BONARDI, C. CORTI, R. CREEMERS, P. A. CROCHET, J. CRNOBRNJA-ISAILOVIC, M. DENOËL, G. F. FICETOLA, J. GONÇALVES, S. KUZMIN, P. LYMBERAKIS, P. DE POUS, A. RODRÍGUEZ, R. SINDACO, J. SPEYBROECK, B. TOXOPEUS, D. R. VIEITES & M. VENCES (2014): Updated distribution and biogeography of amphibians and reptiles of Europe. – *Amphibia-Reptilia*, **35**: 1–31.
- SPITZEN-VAN DER SLUIJS, A., A. MARTEL, J. ASSELBERGHS, E. K. BALES, W. BEUKEMA, M. C. BLETZ, L. DALBECK, E. GOVERSE, A. KERRES, T. KINET, K. KIRST, A. LAUDELOUT, L. F. MARIN DA FONTE, A. NÖLLERT, D. OHLHOFF, J. SABINO-PINTO, B. R. SCHMIDT, J. SPEYBROECK, F. SPIKMANS, S. STEINFARTZ, M. VEITH, M. VENCES, N. WAGNER, F. PASMANS & S. LÖTTERS (2016): Expanding distribution of lethal amphibian fungus *Batrachochytrium salamandrivorans* in Europe. – *Emerging Infectious Diseases*, **22**: 1286–1288.
- SPITZEN-VAN DER SLUIJS, A., F. SPIKMANS, W. BOSMAN, M. DE ZEEUW, T. VAN DER MEIJ, E. GOVERSE, M. KIK, F. PASMANS & A. MARTEL (2013): Rapid enigmatic decline drives the fire salamander (*Salamandra salamandra*) to the edge of extinction in the Netherlands. – *Amphibia-Reptilia*, **34**: 233–239.

- STEGEN, G., F. PASMANS, B. R. SCHMIDT, L. O. ROUFFAER, S. VAN PRAET, M. SCHAUB, S. CANESSA, A. LAUDELLOT, T. KINET, C. ADRIAENSEN, F. HAESEBROUCK, W. BERT, F. BOSSUYT & A. MARTEL (2017): Drivers of salamander extirpation mediated by *Batrachochytrium salamandrivorans*. – *Nature*, **544**: 353–356.
- ŠŤASTNÝ, K., J. ČERVENÝ, J. ROM, M. SOLSKÝ, L. HANEL, J. ANDRESKA, J. VOJAR & K. KEROUŠ (2015): Prague. – pp. 119–153 in: KELCEY, J. G. (ed.): *Vertebrates and Invertebrates of European Cities: Selected Non-Avian Fauna*. – Springer, New York.
- VAN ROOIJ, P., A. MARTEL, F. HAESEBROUCK & F. PASMANS (2015): Amphibian chytridiomycosis: A review with focus on fungus-host interactions. – *Veterinary Research*, **46**: 137.
- VOJAR J., B. HAVLÍKOVÁ, M. SOLSKÝ, D. JABLONSKI, V. IKOVIĆ & V. BALÁŽ (2017): Distribution, prevalence and amphibian hosts of *Batrachochytrium dendrobatidis* in the Balkans. – *Salamandra*, **53**: 44–49.
- YAP, T. A., M. S. KOO, R. F. AMBROSE, D. WAKE & V. T. VREDENBURG (2015): Averting a North American biodiversity crisis. – *Science*, **349**(6247): 481–482.
- ZHU, W., F. XU, C. BAI, X. LIU, S. WANG, X. GAO, S. YAN, X. LI, Z. LIU & Y. LI (2014): A survey for *Batrachochytrium salamandrivorans* in Chinese amphibians. – *Current Zoology*, **60**: 729–735.